

On full-separating sets in graphs

Dipayan Chakraborty^{1,2,3} Annegret Wagler¹

¹Université Clermont-Auvergne, CNRS, Mines de Saint-Étienne,
Clermont-Auvergne-INP, LIMOS, 63000 Clermont-Ferrand, France

²University of Johannesburg, Department of Mathematics and
Applied Mathematics, Auckland Park, 2006, South Africa


³Department of Computer Science and Mathematics,
Lebanese American University, Beirut, Lebanon


CALDAM 2025, PSG College of Technology, Coimbatore, Tamilnadu, India




- 1 Identification problems in graphs : A context for full-separating sets
- 2 Our results
- 3 Questions and future research

Identification problems in graphs: A context for full-separating sets...







Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$


$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$

$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$


Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$

$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

(1) A detector can monitor upto distance 1



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$


Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$

$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:


$$N[v] = N(v) \cup \{v\}$$

$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$

Total-dominating set: A set $C \subseteq V$ such that $N(v) \cap C \neq \emptyset$ for all $v \in V$



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$


$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$

Total-dominating set: A set $C \subseteq V$ such that $N(v) \cap C \neq \emptyset$ for all $v \in V$

(2) A detector can distinguish between itself and a neighbor



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$


$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$

Total-dominating set: A set $C \subseteq V$ such that $N(v) \cap C \neq \emptyset$ for all $v \in V$

(2) A detector can distinguish between itself and a neighbor



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$


$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$

Total-dominating set: A set $C \subseteq V$ such that $N(v) \cap C \neq \emptyset$ for all $v \in V$

(2) A detector can distinguish between itself and a neighbor



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$


$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$

Total-dominating set: A set $C \subseteq V$ such that $N(v) \cap C \neq \emptyset$ for all $v \in V$

(2) A detector can distinguish between itself and a neighbor



Graph $G = (V, E)$

Open neighborhood:

$$N(v) = \{u : uv \in E\}$$

$$N(7) = \{6, 8, 10\}$$

Closed neighborhood:

$$N[v] = N(v) \cup \{v\}$$

$$N[7] = \{6, 8, 10, 7\}$$

“Code” C = set of black vertices

Dominating set: A set $C \subseteq V$ such that $N[v] \cap C \neq \emptyset$ for all $v \in V$

Total-dominating set: A set $C \subseteq V$ such that $N(v) \cap C \neq \emptyset$ for all $v \in V$

Separating set (locating): A set $C \subseteq V$ such that

$$N(u) \cap C \neq N(v) \cap C \iff (N(u) \Delta N(v)) \cap C \neq \emptyset \text{ for all } u, v \in V \setminus C$$

A vertex subset C of a graph G is called...

Locating set: if $(N(u) \Delta N(v)) \cap C \neq \emptyset$ for all $u, v \in V \setminus C$

No faults in detectors: Detectors can distinguish between its vertex and its neighbor
(Slater, 1988)

Closed-separating set: if $(N[u] \Delta N[v]) \cap C \neq \emptyset$ for all $u, v \in V$

Detector fault type 1: Detector cannot distinguish between itself and its neighbors
(Karpovsky et al., 1998)

Open-separating set: if $(N(u) \Delta N(v)) \cap C \neq \emptyset$ for all $u, v \in V$

Detector fault type 2: Detector is completely disabled / destroyed
(Honkala et al., 2002 and Seo et al., 2010)

C. and Wagler, 2024:

Full-separating set: if $(N[u] \Delta N[v]) \cap C = (N(u) \Delta N(v)) \cap C \neq \emptyset$
for all $u, v \in V$

Detector fault type 1 and detector fault type 2

Full-separating dominating code (FD-code):

A set with full-separating property + dominating property

Full-separating total-dominating code (FTD-code):

A set with full-separating property + total-dominating property




Sep	C-Sep		O-Sep		L-Sep		F-Sep	
Code	CD	CTD	OD	OTD	LD	LTD	FD	FTD
adj	$N[u] \triangle N[v]$		$N(u) \triangle N(v)$		$N(u) \triangle N(v)$		$N[u] \triangle N[v]$	
non-adj					$N[u] \triangle N[v]$		$N(u) \triangle N(v)$	
D/TD	$N[u]$	$N(u)$	$N[u]$	$N(u)$	$N[u]$	$N(u)$	$N[u]$	$N(u)$

Existence of full-separating sets:

The graph must be twin-free = open-twin-free + closed-twin-free


$X \in \text{CODES} = \{\text{LD, LTD, ID, ITD, OD, OTD, FD, FTD}\}$

X-number of a graph G : $\gamma^X(G) = \min\{|C| : C \text{ is an X-code of } G\}$



Our results...

Relations to other codes...



$X' \longrightarrow X$ stands for $\gamma^{X'}(G) \leq \gamma^X(G)$

An interesting result...

Theorem (C. and Wagler, 2024)

For a twin-free and isolate-free graph $G = (V, E)$, we have

$$\gamma^{\text{FTD}}(G) - 1 \leq \gamma^{\text{FD}}(G) \leq \gamma^{\text{FTD}}(G).$$

Proof sketch.

Clearly, $\gamma^{\text{FD}}(G) \leq \gamma^{\text{FTD}}(G)$ since any FTD-code is also an FD-code.

To prove the other inequality, take a minimum FD-code C of G .

Notice that there exists at most one $v \in V$ such that $N(v) \cap C = \emptyset$. Thus C is “almost” total-dominating.

To turn C into an FTD-code, we may have to include in C one neighbor of v . This implies $\gamma^{\text{FTD}}(G) \leq |C| + 1 = \gamma^{\text{FD}}(G) + 1$.

Computational complexities...

FD-CODE

Input: (G, k) : A graph G and a positive integer k .

Problem: Does there exist an FD-code C of G with $|C| \leq k$?

FTD-CODE

Input: (G, k) : A graph G and a positive integer k .

Problem: Does there exist an FTD-code C of G with $|C| \leq k$?

FD = FTD - 1

Input: A graph G and an integer k .

Problem: Is $\gamma^{\text{FTD}}(G) = k$ and $\gamma^{\text{FD}}(G) = k - 1$?

Computational complexities...

FD-CODE

Input: (G, k) : A graph G and a positive integer k .

NP-hard!

Problem: Does there exist an FD-code C of G with $|C| \leq k$?

FTD-CODE

Input: (G, k) : A graph G and a positive integer k .

NP-hard!

Problem: Does there exist an FTD-code C of G with $|C| \leq k$?

FD = FTD - 1

Input: A graph G and an integer k .

NP-hard!

Problem: Is $\gamma^{\text{FTD}}(G) = k$ and $\gamma^{\text{FD}}(G) = k - 1$?


Theorem (C. and Wagler, 2024)

FTD-CODE is NP-complete.

Proof sketch.

Reduction from 3-SAT with formula ψ on n variables and m clauses.

E.g. $\psi = (x \vee \neg y \vee z) \wedge (\neg x \vee \neg z \vee w) \wedge (\neg y \vee z \vee \neg w)$.



Theorem (C. and Wagler, 2024)


FTD-CODE is NP-complete.

Proof sketch.

Reduction from 3-SAT with formula ψ on n variables and m clauses.

E.g. $\psi = (x \vee \neg y \vee z) \wedge (\neg x \vee \neg z \vee w) \wedge (\neg y \vee z \vee \neg w)$.

ψ satisfiable $\iff (G^\psi, k = 7n + 2m)$ is YES-instance of FTD-CODE



Theorem (C. and Wagler, 2024)


FTD-CODE is NP-complete.

Proof sketch.

Reduction from 3-SAT with formula ψ on n variables and m clauses.
E.g. $\psi = (x \vee \neg y \vee z) \wedge (\neg x \vee \neg z \vee w) \wedge (\neg y \vee z \vee \neg w)$.

ψ satisfiable $\iff (G^\psi, k = 7n + 2m)$ is YES-instance of FTD-CODE

ψ satisfiable $\implies \exists C$ such that $|C| = \gamma^{\text{FTD}}(G^\psi) = k$



Theorem (C. and Wagler, 2024)


FTD-CODE is NP-complete.

Proof sketch.

Reduction from 3-SAT with formula ψ on n variables and m clauses.
E.g. $\psi = (x \vee \neg y \vee z) \wedge (\neg x \vee \neg z \vee w) \wedge (\neg y \vee z \vee \neg w)$.

ψ satisfiable $\iff (G^\psi, k = 7n + 2m)$ is YES-instance of FTD-CODE

ψ satisfiable $\iff \exists C$ such that $|C| = \gamma^{\text{FTD}}(G^\psi) = k$



FTD-CODE is NP-complete.


Proof sketch.

Reduction from 3-SAT with formula ψ on n variables and m clauses.
 E.g. $\psi = (x \vee \neg y \vee z) \wedge (\neg x \vee \neg z \vee w) \wedge (\neg y \vee z \vee \neg w)$.

ψ satisfiable $\iff (G^\psi, k = 7n + 2m)$ is YES-instance of FTD-CODE

ψ satisfiable $\iff \exists C$ such that $|C| = \gamma^{\text{FTD}}(G^\psi) = k$


ψ satisfiable $\iff (G^\psi, 7n + 2m - 1)$ is YES-instance of FD-CODE



Theorem (C. and Wagler, 2024)

Let G be either a path P_n for $n \geq 4$ or a cycle C_n for $n \geq 5$. Moreover, let $n = 6q + r$ for non-negative integers q and $r \in [0, 5]$. Then

$$\gamma^{\text{FD}}(G) = \gamma^{\text{FTD}}(G) = \begin{cases} 4q + r, & \text{if } r \in [0, 4]; \\ 4q + 4, & \text{if } r = 5. \end{cases}$$




Results on some other selected graph classes...

A subclass of bipartite graphs:

For a half-graph H_k with $k \geq 3$, we have

$$\gamma^{\text{FTD}}(H_k) = 2k \quad \text{and} \quad \gamma^{\text{FD}}(H_k) = 2k - 1.$$




H_4

A subclass of split graphs:

For a thin headless spider S_k with $k \geq 4$, we have

$$\gamma^{\text{FTD}}(S_k) = 2k - 1 \quad \text{and} \quad \gamma^{\text{FD}}(S_k) = 2k - 2.$$




S_4

A subclass of split graphs:

For a thick headless spider \bar{S}_k with $k \geq 4$, we have

$$\gamma^{\text{FTD}}(\bar{S}_k) = 2k - 2 = \gamma^{\text{FD}}(\bar{S}_k).$$



\bar{S}_4

Questions and future research...

Questions and future research...

- Study bounds and other combinatorial aspects of the FD- and FTD-codes on other well-known graphs classes.
- Study the new code numbers ($\gamma^{\text{FD}}(\cdot)$ and $\gamma^{\text{FTD}}(\cdot)$) with respect to other previously introduced code numbers like $\gamma^{\text{LD}}(\cdot)$, $\gamma^{\text{OTD}}(\cdot)$ etc.
- Explore the algorithmic aspects of FD-CODE and FTD-CODE more.

Thank you!